



Identification of steroid compound using preparative Thin Layer Chromatography, GC-MS and antimicrobial and antioxidant properties of *Cenchrus setigerus* (Poaceae)

Premlata Singariya^{1*}, Padma Kumar¹ and Krishan Kumar Mourya²

1, Dr. D.S. Kothari Post Doctoral Fellow, Laboratory of Tissue Culture and Secondary Metabolites,
Department of Botany, University of Rajasthan, Jaipur, (R.J.) - India

2, Veterinary Officer & In-charge, 1st Gr. Govt. Veterinary Hospital, Pahari, Bharatpur, (R.J.) - India

Abstract

The present investigation was carried out to determine the possible bioactive components of *Cenchrus setigerus* using PTLC, IR spectra and GC-MS analysis, *in vitro* antibacterial activity of methanolic extracts and comparative estimation of antioxidant properties (*in-vitro* and *in-vivo*) in terms of POX, CAT, PPO, SOD, LPO activity and carotenoids, total phenolics contents of *C. setigerus* (Poaceae) were done. Antibacterial activity was evaluated against seven Gram-negative bacteria, two Gram-positive bacteria and three fungi, using disk diffusion method followed by determination of minimum inhibitory concentrations (MIC) by broth dilution method, against sensitive bacteria and fungi. Result reveals that the most bioactive compound in *C. setigerus* was stigmasterol. The highest activity was exhibited by the root extract against *Bacillus subtilis*. *Klebsiella pneumoniae*, *Proteus mirabilis* and *Aspergillus flavus* were the most sensitive pathogens which show maximum antimicrobial effects. Highest antioxidant properties of *C. setigerus* were found to be greater in *in-vitro* than *in-vivo*.

Key-Words: Antibacterial activity, stigmasterol, Minimum Inhibitory Concentration (MIC), GC-MS, PTLC, *Cenchrus* grass.

Introduction

The chemical analysis of methanolic extract of *Cenchrus setigerus* showed a mixture of long-chain hydrocarbons, carboxyl esters, alcohols, acids, alkaloids, steroids, amino and nitro compound etc. Stigmasta-5,22-dien-3-ol (stigmasterol) was isolated and identified from the extract. Phytochemical screening using the pharmacognostic methods revealed the presence of flavonoids, steroids and alkaloids. Taking into consideration of the medicinal importance of this plant, the methanolic extract of *C. setigerus* was analyzed for the first time using GC-MS. This work will help to identify the compounds of therapeutic value. GC-MS is the best technique to identify the bioactive constituents of long chain hydrocarbons, alcohols, acids, ester, alkaloids, steroids, amino and nitro compound etc.

Reactive oxygen species (ROS) are inevitable byproducts from the essential aerobic metabolisms (caused oxidative damages). It causes peroxidation of polyunsaturated fatty acids in the membranes (Smirnoff, 1995) and they need to be maintained under sub-lethal levels for normal plant growth. Hence, plants are equipped with an array of enzymatic and non-enzymatic antioxidant molecules to alleviate cellular damage caused by ROS (Apel and Hirt, 2004). Multiple antioxidant enzymes systems are involved in the enzymatic scavenging of ROS. Superoxide dismutases (SOD) react with the superoxide radical to produce H₂O₂. Hydrogen peroxide is scavenged by catalases (CAT) and peroxidases (POX).

In the present scenario development of antimicrobial resistance to anti microbial agents has led to treatment failure and the shift of medical care from orthodox to herbal medicine. Various plant extracts can serve both as potential antimicrobial crude drugs as well as a source of new anti-infective agents (Rios, 2005).

Cenchrus setigerus (C₄ grass) is gaining attention in various field of research, as they are best suited to the present environmental conditions. This grass is more

*** Corresponding Author**

E.mail:premlatasingariya@gmail.com,
mourya07@gmail.com

efficient at gathering Carbon dioxide and utilizing nitrogen from the atmosphere and recycled N in the soil (Bessman, 1956; Singariya, 2009). It has excellent soil binding capacity which helps to conserve soil in desert areas (Sinha *et al.*, 1996) and more competitive under the conditions of high temperature, solar radiation and low moisture (Agrawal, 2007).

Material and Methods

Experimental design: [1] Identification of Bioactive Components:

(A) Identification of Components by GC-MS: Interpretation on mass spectrum of GC-MS was done using the database of National Institute of standard and Technology (NIST) having more than 62,000 patterns. The mass spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library. The name, molecular weight and structure of the components of the test materials were ascertained.

(B) Separation of active compound from *Cenchrus setigerus* extracts suspension by preparative thin layer chromatography (PTLC): The glass plates (20 x 20 cm) coated with silica gel 'G' (0.4-0.5 mm thick and 45 gm/80 ml distilled water) were dried at room temperature (Meena and Patni, 2008). The dried plates were activated at 100°C for 30 minutes in an oven and cooled at room temperature. Methanolic extracts were separately applied 1 cm above the edge of the plates along with standard reference compound (Stegmasterol).

(C) Development of chromatogram: The development tank was saturated with suitable solvent system hexane, acetone (8:2), benzene, ethyl acetate (85:15 or 3:1), chloroform, methanol and water (10:10:3) for the analysis of lipid present in plant extract but, hexane, acetone (8:2) gave better separation (Kaul and Staba, 1968).

(D) Spot visualization and identification by TLC and IR spectra: Few pieces of iodine crystals were kept in the tank and covered with glass plate to saturate the tank with iodine vapor. The plate was then kept in iodine vapor saturated tank and left for few hours. Melting point and IR spectra of each of the isolated compound was taken and a comparison of TLC color reaction was made, which was found to be in accordance with those reported for authentic compounds.

[2] For Antimicrobial Activity: (i) Plant material: *C. setigerus* were collected in the month of August from the CAZRI, Jodhpur (Rajasthan). The collected plant materials were transferred immediately to the laboratory cleaned with water and selected plant parts were separately shade dried until weight has been

constant. Shade dried parts were powdered with the help of grinder (Singariya *et al.*, 2012 l).

(ii) Preparation of extracts: Plant samples (root, stem, leaf and seeds) were extracted by the hot extraction technique (Harborne, 1984) by using the soxhlet apparatus for 18 hours at a temperature not exceeding the boiling point of methyl alcohol (65°C) following the method of Subramanian and Nagarjan, (1969). The obtained extracts were filtered by using Whatman No. 1 filter paper and then concentrated at 40°C by using an evaporator. The residual extracts were stored in refrigerator at 4°C in small and sterile glass bottles for antimicrobial activity and the crude extract was dissolved in chloroform before chromatographic examination (Rishi *et al.*, 2011).

(iii) Micro-organisms: The organisms used in this study were seven Gram-negative bacteria, two Gram-positive bacteria and three fungi (table 1). Selected microorganisms were procured from IMTECH, Chandigarh, India.

(iv) Preparation of test pathogens and Disc diffusion assay: Bacterial strains were grown and maintained on NA medium, while fungi were maintained on SDA medium. DDA was performed for screening by standard method (Singariya, *et al.*, 2012h). AI for each extract was calculated.

$$\text{Activity index (AI)} = \frac{\text{Inhibition Zone of the sample}}{\text{Inhibition Zone of the standard}}$$

(v) Serial dilution method: MIC was determined as the least extract concentration which inhibited the growth of the test organisms (Jain and Sharma, 2009; Singariya, *et al.*, 2012g). Bacterial and fungal suspensions were used as negative control, while broth containing standard drug was used as positive control.

(vi) Determination of Minimum bactericidal/fungicidal concentration (MBC/MFC): Equal volume of the various concentration of each extract and nutrient broth mixed in micro-tubes to make up 0.5ml of solution. 0.5ml of McFarland standard of the organism suspension was added to each tube (Bhattacharya, *et al.*, 2009). The tubes were incubated aerobically and MBC was determined by sub culturing and further incubated for 24 h. The highest dilution that yielded no single bacterial colony was taken as the MBC (Akinyemi *et al.*, 2005).

(vii) Total activity (TA) determination: Total activity is the volume at which the test extract can be diluted with the ability to kill microorganisms. It is calculated by dividing the amount of extract from 1 g plant material by the MIC of the same extract or compound

isolated and is expressed in ml/g (Singariya, *et al.*, 2012f).

Extract per gram dried plant part

Total Activity = -----

MIC of extract

[3] For Antioxidant activity: (a) *In vivo* studies: For *in vivo* studies seeds of *C. setigerus* were grown in 12" earthenware pots. Pots were filled with 8kg of a mixture of garden soil and goat manure in the ratio of 3:1. Pots were watered everyday. After two weeks of sowing thinning was done and 3-4 plants of uniform size were selected in each pot. Leaf samples were collected for biochemical analysis (Singariya *et al.*, 2009).

(b) *In vitro* studies: For *in vitro* studies seeds of *C. setigerus* were first sterilized by treating them with 0.01% HgCl₂ solution for five minutes, these sterilized seeds were then germinated in 9cm Petri plates lined with double layer of filter paper in distilled water. These Petri plates were kept in BOD incubator at 28°C±2°C. A cool fluorescent light of 34.1µmol/m²/s PAR was given to the seeds. Five-day-old seedlings were analyzed for different parameter (Chouhan *et al.*, 2007).

(c) Preparation of enzyme extract: All operations for preparing the enzyme extracts were performed at 4°C. Plant material (*in vitro*-seedling, *in vivo*-leaves) were homogenized using appropriate buffer in pre-chilled pestle mortar and centrifuged at 10,000 rpm for 20 min. The supernatant collected was used for all the enzyme assays and metabolites estimation.

(d) Estimation of Antioxidant: The level of lipid peroxidation (LPO) was measured in terms of malondialdehyde (MDA) content a product of lipid peroxidation described by Heath and Packer (1968). The activity of catalase (CAT, EC.1.11.1.6), peroxidase (POX, EC.1.11.1.7) and polyphenol oxidase (PPO, EC.1.10.3.1) was assayed after the method of Chance and Maehly (1955) with the some modification. Superoxide dismutase (SOD, EC.1.15.1.1) activity was estimated by the method of Beauchamp and Fridovich (1971). The total phenolics in extracts determined according to Folin-Ciocalteu procedure (Singleton and Rossi, 1995). Arnon's (1949) method was used for calculation of carotenoids content.

Results and Discussion

[1] Identification of Bioactive Components: TLC plate of *Cenchrus setigerus* extracts were visualized under UV lamp one of the spot gave characteristic fluorescence and their R_f values were comparable to their respective standard compound (Stigmasterol-grayish violet, R_f value- 0.83) (Fig.-1). The

characteristic color was also developed when TLC plates were sprayed with anisaldehyde reagent (Stigmasterol- grayish violet) corresponding to their authentic standard compound. Melting point (Stigmasterol 131-132°C) was also measured and compared with authentic standard compound.

GC- MS analysis: GC – MS was done using the database of National Institute of standard and Technology (NIST) (Fig.-2). The GC-MS studied showed that the retention time and peaks of the isolated Stigmasterol was comparable with that of standard. The active principles with their retention time (34.342) and concentration (4% area) in the standard and sample are presented. However, till date there was no report on the presence of sterols from *C. setigerus*. In the present study Stigmasterol has been confirmed in *C. setigerus* of family Poaceae.

[2] Antimicrobial Assay: Antimicrobial assay (assessed in terms of ZOI, AI, MIC, MBC/MFC and TA) (table 2, 3, 4) of the Methanolic extracts of *C. setigerus* against selected microorganisms were recorded. The antibacterial activity of the test samples was determined by measuring the diameter of zone of inhibition expressed in mm. The assay was repeated twice and mean of the three experiments was recorded.

(A) Root: Root extract show highest activity ZOI-12.50±0.64 mm, AI-0.625 followed by ZOI-12.33±0.26 mm, AI-0.617 against *B. subtilis* and *K. pneumoniae* respectively. Lowest MIC value 0.469 mg/ml and highest total activity value 70.701 were recorded against *Agerobacterium tumefaciens*.

(B) Stem: Stem extract show highest activity ZOI-12.33±0.24 mm, AI-1.028 followed by ZOI-11.67±0.24 mm, AI-0.584 against *P. merabilis* and *K. pneumoniae* respectively. Lowest MIC value 0.938 mg/ml and highest total activity value 40.533 were recorded against *A. tumefaciens*.

(C) Leaf: Leaf extract show highest activity ZOI-9.50±0.64 mm, AI-0.792 followed by ZOI-9.33±0.25 mm, AI-1.166 against *P. merabilis* and *Aspergillus flavus* respectively. Lowest MIC values were recorded against *Escherichia coli*, *B. subtilis*, *K. pneumoniae* and *A. flavus*.

(D) Seed: Seed extract show highest activity ZOI-10.67±0.22 mm, AI-0.889 followed by ZOI-9.67±0.21 mm, AI-0.484 against *P. merabilis* and *K. pneumoniae* respectively. Lowest MIC value 1.875 mg/ml and highest total activity value 16.107 were recorded against *A. tumefaciens*.

Phyto-chemical estimation: The phyto-chemical estimation for the *C. setigerus* was carried out according to Farnsworth (1966) wherein the consistency was found to be sticky in all the extracts.

The yield of the extracts was also analyzed where in the highest yields were recorded for leaf extract of *C. setigerus* (4.06%) (table 5)

Most susceptible organisms in the investigation were *P. merabilis* and *K. pneumoniae* against which, all the plant extracts showed zone of inhibition. But, according to the zone of inhibition *B. subtilis* was the most susceptible organism, which supported the finding that plant extracts are usually more active against Gram positive bacteria than Gram negative (Singariya, *et al.*, 2011a, b, c, d, 2012a; Lin *et al.*, 1999; Palombo and Semple, 2001). The Gram-negative bacterial cell wall outer membrane appears to act as a barrier to many substances including synthetic and natural antibiotics (Tortora *et al.*, 2001). *A. tumefaciens* was the most sensitive bacteria according to lowest MIC value and highest total activity value (Singariya, *et al.*, 2012b, c). Previous studies have noted alcohols to be reliable and consistent solvents for the extraction of antimicrobial substances from medicinal plants (Ahmad *et al.*, 1998).

[3] Antioxidant activity: In the present study, comparative estimation of antioxidant properties (*In-vitro* and *in-vivo*) of *C. setigerus* in terms of POX, CAT, PPO, SOD, LPO activity and carotenoids, total phenolics contents were done. POX, CAT, SOD, LPO and PPO activity were found to greater in *in-vitro* than *in-vivo* estimation (table 6) but, *visé versa* in case of carotenoids and total phenolics contents (graph 1). As a part of the enzymatic component of antioxidant system of plant response, antioxidant enzymes contribute significantly to ROS detoxification. The coordinate function of antioxidant enzymes such as SOD, POX, CAT and GR helps in processing of ROS and regeneration of redox ascorbate and glutathione metabolites (Almeselmani *et al.*, 2006). Abiotic stress (water, salinity, Heat, Heavy metal) induces generation of active oxygen species (Cadenas, 1989) causing lipid peroxidation and consequently membrane injury, protein degradation, enzymes inactivation, pigment bleaching and disruption of DNA strands (Deshmukh and Dhupal, 2005).

Acknowledgement

Authors are expressing their thanks to UGC for providing the funds for this project under Dr. D. S. Kothari, Post doctoral fellowship scheme.

References

1. Agrawal P. (2007). *Ecophysiological and Biochemical studies Related to drought adaptation in grasses of Indian Desert*. Ph.D. Thesis, J. N. Vyas University.
2. Ahmad I., Mehmood Z. and Mohammad F. (1998). Screening of some Indian medicinal plants for

their antimicrobial properties. *J Ethnopharmacol* 62:183-193.

3. Akinyemi K.O., Oladapo O., Okwara C.E., Ibe C.C. and Fasura K.A. (2005). Screening of crude extracts of six medicinal plants used in southwest Nigerian unorthodox medicine for antimethicillin resistant *Staphylococcus aureus* activity. *BMC Complementary and Alternative Medicine* 5: 6.
4. Almeselmani M., Deshmukh R.K., Sairam R.K., Kushwaha S.R. and Singh T.P. (2006). Protective role of antioxidant enzymes under high temperature stress. *Plant Sci*, 171: 382-388.
5. Apel K. and Hirt H. (2004). Reactive oxygen species: metabolism, oxidative stress and signal transduction. *Annu. Rev. Plant Biol.* 55: 373-399.
6. Arnon D.I. (1949). Copper enzymes in isolated chloroplast: polyphenol oxidase in *Beta vulgaris*. *Plant Physiol*, 24: 1-5.
7. Beauchamp C. and Fridovich I. (1971). Superoxide dismutase: Improved assay and an assay applicable to acrylamide gels. *Anal. Biochem* 44: 276-287.
8. Bessman S.P. (1956). *Ammonia Metabolism in Animals: Symposium on Inorganic Nitrogen Metabolism*. Mc Elry and Glass (eds.) The Johns Hopkins Press.
9. Bhattacharya S., Zaman M.K., Haldar P.K. (2009). Antibacterial Activity of stem Bark and Root of Indian *Zanthoxylum nitidum*, *Asian J. of Pharm. and Clin. Res.*, 2(1): 30-34.
10. Cadenas E. (1989). Biochemistry of oxygen toxicity. *Annu. Rev. Biochem.* 58: 79-110.
11. Chance B. and Maehly A.C. (1955). Assay of catalases and peroxidases. *Methods Enzymol*, 2: 764-775.
12. Chouhan J.K., Rathi P., Singariya P. and Bohra S.P. (2007). Effect of Salicylic Acid in Response to Applied Moisture Stress on Nitrogen Assimilation in *Lablab* Bean. *Journal of Current Agriculture*, 31(1-2): 79-83.
13. Deshmukh R.N. and Dhaumal K.N. (2005). Enzymatic studies in *Sorghum* cultivars under PEG induced water stress. *Indian J. Plant Physiol.*, 10(4): 349-353.
14. Farnsworth N.R. (1966). Biology and Phytochemical screening of plants. *Pharm Sci.*, 55: 225-276.
15. Harborne J.B. (1984). *Phytochemical Methods*, Chapman and Hall, London, New York p.p. 5-6.
16. Heath R.L. and Packer L. (1968). Photo peroxidation in isolated chloroplast. *Arch. Biochem. Biophys.*, 125: 189-198.

17. Jain T. and Sharma K. (2009). Assay of antibacterial activity of *Polyalthia longifolia* Benth. And Hook. Leaf extracts. *Journal of Cell and Tissue Research*, 9(2): 1817-1820.
18. Kaul B. and Staba E.J. (1968). *Dioscorea* tissue cultures. I. Biosynthesis and isolation of diosgenin from *Dioscorea deltoidea* callus and suspension cells. *Lloydia*, 31: 171-179.
19. Lin J., Opaque A.R., Geheeb-Keller M., Hutchings A.D., Terblanche S.E. and Jäger A.K. (1999). Preliminary screening of some traditional Zulu medicinal plants for anti-inflammatory and antimicrobial activities. *J Ethnopharmacol.*, 68: 267-274.
20. Meena M.C. and Patni P. (2008). Isolation and Identification of Flavonoid "Quercetin" from *Citrullus colocynthis* (Linn.) Schrad. *Asian J. Exp. Sci.*, 22(1): 137-142.
21. Palombo E.A. and Semple S.J. (2001). Antibacterial activity of traditional Australian medicinal plants. *J Ethnopharmacol.*, 77: 151-157.
22. Rios J.L. and Reico M.C. (2005). Medicinal plants and antimicrobial activity. *J Ethnopharmacol.*, 100: 80-84.
23. Rishi A., Sharma A. and Meena S. (2011). Quantitative Estimation of Beta-Sitosterol and Stigmasterol in-vivo and in-vitro *Terminalia chebula* Ritz. *I. Res. j of Pharmacy*, 2(3): 115-116.
24. Singariya P. (2009). *Effect of Sub-Optimal Environment and PGR's on Metabolic Pattern of Certain Species of Cenchrus*. Ph.D Thesis, J. N. Vyas University.
25. Singariya P., Kumar P. and Mourya K.K. (2012a). Phyto-chemical Screening and Antimicrobial Activities of Dhaman grass and Indian Ginseng. *Journal of Pharmacy Res.*, 5(1): 135-139.
26. Singariya P., Kumar P. and Mourya K.K. (2012b). Evaluation of Antibacterial Activity and Preliminary Phytochemical Studies on the Stem of *Cenchrus ciliaris* and *Cenchrus setigerus*. *Asian J Pharm Clin Res.*, 5(1): 163-167.
27. Singariya P., Kumar P. and Mourya K.K. (2012c). Antimicrobial Activity of Fruit coat (Calyx) of *Withania somnifera* against some Multi Drug Resistant Microbes *International Journal of Biological & Pharmaceutical Research*, 3(2): 252-258.
28. Singariya P., Kumar P. and Mourya K.K. (2012f). Bio-efficacy of Unripen Fruits of Winter Cherry (*Withania somnifera*). *Res. J. of Pharm., Biol. and Chemical Sci.*, 3(2): 479-487.
29. Singariya P., Kumar P. and Mourya K.K. (2012g). Ripen Fruits of Indian Ginseng: Phyto-chemical and Pharmacological examination against Human and Plant Pathogens. *International Journal of Applied Biology and Pharmaceutical Technology*, 3(2): 1-8.
30. Singariya P., Kumar P. and Mourya K.K. (2012h). Comparative Pharmacological Study of Fruits and Flowers Extract of *Withania somnifera*. *International Journal of Pharmaceutical and Phyto-pharmacological Research*, 1(5): 276-282.
31. Singariya P., Kumar P. and Mourya K.K. (2012l). Estimation of Bio-activity of Aerial parts of *Withania somnifera* Against the Bacterial and Fungal Microbes. *International Journal of Pharmacy and Pharmaceutical science*, 4(3): 553-557.
32. Singariya P., Mourya K.K. and Kumar P. (2011a). Preliminary Phyto-profile and Pharmacological Evaluation of some Extracts of *Cenchrus* grass against Selected Pathogens. *J. Pharm. Sci. and Res.*, 3(8): 1387-1393.
33. Singariya P., Mourya K.K. and Kumar P. (2011b). Bio Activity of crude extracts of Leaves of *Cenchrus* Grass in different polar solvents against some pathogenic microbes. *I. J. of Pharm. Sci. Review and Res.*, 11(1): 124-129.
34. Singariya P., Mourya K.K. and Kumar P. (2011c). Comparative Microcidal Activity of *Withania somnifera* and *Cenchrus setigerus* against the Pathogenic Micro-organisms. *I. J. of Pharmacy and Pharm. Science*, 3(55): 511-515.
35. Singariya P., Mourya K.K. and Kumar P. (2012d). Antimicrobial Activity of the Crude Extracts of *Withania somnifera* and *Cenchrus setigerus* *In-vitro*. *Pharmacognosy journal*, 3(27): 60-65.
36. Singariya P., Rathi P., Gadi B.R. and Bohra S.P. (2009). Effect of Ethephon on the Activities of Ammonia Assimilation Enzymes and Metabolites of *Cenchrus Setigerus*-76 grown *in vivo*. *Advances in Microbiology & Biotechnology*, p.p. 347-351.
37. Singleton S. and Rossi T. (1995). *Methodology of plant metabolites and enzymes*. Vol. II, (eds.) Bhansali RR, Jindal SK, pp. 245-265.
38. Sinha R.K., Bhatia S. and Vishnoi R. (1996). *Desertification control and rangeland management in the Thar desert of India*. In: Rala Report no. 200, pp. 115-123.
39. Smirnoff N. (1995). Antioxidant systems and plant response to the environment. In "*Environment and Plant Metabolism: Flexibility and Acclimation*" (N Smirnoff ed), Bios Scientific, Oxford, pp. 217-243.
40. Subramanian S.S. and Nagarjan S. (1969). Flavonoids of the seeds of *Crotolaria retusa* and *Crotolaria striata*. *Current Sci.*, 38: 65.

41. Tortora G.J., Funke B.R. and Case C.L. (2001).
Microbiology: An Introduction, Benjamin
Cummings. San Francisco.

Table 1: Name of the tested pathogens

| S/No. | Pathogens | Name of Pathogens | G+ve/ G-ve | Specimen no. |
|-------|-----------|-----------------------------------|------------|--------------|
| 1. | Bacteria | <i>Escherichia coli</i> | G-ve | MTCC-46 |
| 2. | | <i>Staphylococcus aureus</i> | G+ve | MTCC-3160 |
| 3. | | <i>Raoultella planticola</i> | G-ve | MTCC-530 |
| 4. | | <i>Pseudomonas aeruginosa</i> | G-ve | MTCC-1934 |
| 5. | | <i>Bacillus subtilis</i> | G+ve | MTCC-121 |
| 6. | | <i>Enterobacter aerogens</i> | G-ve | MTCC-111 |
| 7. | | <i>Proteus merabilis</i> | G-ve | MTCC-530 |
| 8. | | <i>Klebsiella pneumoniae</i> | G-ve | MTCC-3310 |
| 9. | | <i>Agerobacterium tumefaciens</i> | G-ve | MTCC-431 |
| 10. | Fungi | <i>Candida albicans</i> | - | MTCC-183 |
| 11. | | <i>Aspergillus flavus</i> | - | MTCC-277 |
| 12. | | <i>Aspergillus niger</i> | - | MTCC-282 |

Table 2: Zone of Inhibition (mm)* and Activity index of Methanolic extract *C. setigerus*

| Test Pathogens | Bio-activity of different parts of <i>Cenchrus setigerus</i> against pathogens | | | | | | | |
|-----------------------|--|-------|------------|-------|-----------|-------|------------|-------|
| | Root | | Stem | | Leaf | | Seed | |
| | ZOI | AI | ZOI | AI | ZOI | AI | ZOI | AI |
| <i>E. coli</i> | 11.17±0.24 | 0.559 | 10.67±0.28 | 0.534 | 8.50±0.64 | 0.425 | 9.33±0.24 | 0.467 |
| <i>S. aureus</i> | - | - | - | - | - | - | - | - |
| <i>R. planticola</i> | - | - | - | - | - | - | - | - |
| <i>P. aeruginosa</i> | - | - | - | - | - | - | - | - |
| <i>B. subtilis</i> | 12.5±0.64 | 0.625 | - | - | 9.33±0.24 | 0.467 | - | - |
| <i>P. merabilis</i> | 12.17±0.29 | 1.014 | 12.33±0.24 | 1.028 | 9.50±0.64 | 0.792 | 10.67±0.22 | 0.889 |
| <i>E. aerogens</i> | - | - | - | - | - | - | - | - |
| <i>K. Pnemoniae</i> | 12.33±0.26 | 0.617 | 11.67±0.24 | 0.584 | 8.5±0.64 | 0.425 | 9.67±0.21 | 0.484 |
| <i>A. tumefaciens</i> | 11.67±0.28 | 0.729 | 10.5±0.64 | 0.656 | 8.17±0.24 | 0.511 | 9.33±0.22 | 0.583 |
| <i>C. albicans</i> | - | - | - | - | - | - | - | - |
| <i>A. flavus</i> | 8.17±0.26 | 1.021 | - | - | 9.33±0.25 | 1.166 | 8.33±0.22 | 1.041 |
| <i>A. niger</i> | - | - | - | - | - | - | - | - |

*All values are mean ± SD, n=3, ZOI-Zone of Inhibition, AI-Activity Index

Table 3: MIC and MBC/MFC by methanolic extract of *C. setigerus* against tested pathogens

| Test Pathogens | Bio-activity of different parts of <i>Cenchrus setigerus</i> against pathogens | | | | | | | |
|-----------------------|--|---------|-------|---------|------|---------|-------|---------|
| | Root | | Stem | | Leaf | | Seed | |
| | MIC | MBC/MFC | MIC | MBC/MFC | MIC | MBC/MFC | MIC | MBC/MFC |
| <i>E. coli</i> | 1.875 | 3.75 | 3.75 | 7.5 | 3.75 | 7.5 | 3.75 | 3.75 |
| <i>S. aureus</i> | - | - | - | - | - | - | - | - |
| <i>R. planticola</i> | - | - | - | - | - | - | - | - |
| <i>P. aeruginosa</i> | - | - | - | - | - | - | - | - |
| <i>B. subtilis</i> | 1.875 | 3.75 | - | - | 3.75 | 7.5 | - | - |
| <i>P. merabilis</i> | 3.75 | 7.5 | 3.75 | 3.75 | 7.5 | 15 | 3.75 | 7.5 |
| <i>E. aerogens</i> | - | - | - | - | - | - | - | - |
| <i>K. Pnemoniae</i> | 1.875 | 1.875 | 1.875 | 3.75 | 3.75 | 7.5 | 3.75 | 3.75 |
| <i>A. tumefaciens</i> | 0.469 | 0.938 | 0.938 | 1.875 | 7.5 | 7.5 | 1.875 | 1.875 |
| <i>C. albicans</i> | - | - | - | - | - | - | - | - |
| <i>A. flavus</i> | 7.5 | 15 | - | - | 3.75 | 7.5 | 3.75 | 7.5 |
| <i>A. niger</i> | - | - | - | - | - | - | - | - |

Table 4: Total activity of methanolic extract of different parts of *C. setigerus*

| S/No. | Test Pathogens | Total activity of different parts | | | |
|-------|-----------------------|-----------------------------------|--------|--------|--------|
| | | Root | Stem | Leaf | Seed |
| 1. | <i>E. coli</i> | 19.893 | 10.133 | 10.827 | 8.053 |
| 2. | <i>S. aureus</i> | - | - | - | - |
| 3. | <i>R. planticola</i> | - | - | - | - |
| 4. | <i>P. aeruginosa</i> | - | - | - | - |
| 5. | <i>B. subtilis</i> | 19.893 | - | 10.827 | - |
| 6. | <i>P. merabilis</i> | 9.947 | 10.133 | 5.413 | 8.053 |
| 7. | <i>E. aerogens</i> | - | - | - | - |
| 8. | <i>K. Pnemoniae</i> | 19.893 | 20.267 | 10.827 | 8.053 |
| 9. | <i>A. tumefaciens</i> | 79.701 | 40.533 | 5.413 | 16.107 |
| 10. | <i>C. albicans</i> | - | - | - | - |
| 11. | <i>A. flavus</i> | 4.973 | - | 10.827 | 8.053 |
| 12. | <i>A. niger</i> | - | - | - | - |

Table 5: Phyto-profile of Methanolic extracts of different parts of *C. setigerus*

| S/No. | Parts | % Yield | Color | Consistency |
|-------|-------|---------|------------|-------------|
| 1. | Root | 3.73 | Brown | Sticky |
| 2. | Stem | 3.80 | Dark brown | Sticky |
| 3. | Leaf | 4.06 | Dark green | Sticky |
| 4. | Seed | 3.02 | Brown | Sticky |

Table 6: Antioxidants of *Cenchrus setigerus* (*In-vitro* and *In-vivo*)

| S/No. | Antioxidants | Parameter | (<i>In-vitro</i>) | (<i>In-vivo</i>) |
|-------|-----------------|--|---------------------------|--------------------|
| 1. | POX | (μ mol H ₂ O ₂ /min/g F. wt.) | 0.742±0.045 | 0.542±0.035 |
| 2. | CAT | (Units/min/g F. wt.) | 1.503±0.016 | 1.34±0.012 |
| 3. | PPO | (min ⁻¹ g ⁻¹ F. wt.) | 2.92±0.032 | 2.27±0.032 |
| 4. | SOD | (μ mol nitro blue toluene /sec./g D. wt.) | 0.487±0.036 | 0.261±0.021 |
| 5. | LPO | (m mol MDA/g F. wt.) | 7.283±0.126 | 6.83±0.16 |
| 6. | Carotenoids | (mg/g F. wt.) | 0.063±0.007 ^{NS} | 0.159±0.008 |
| 7. | Total Phenolics | (mg/g F. wt.) | 11.35±0.135 | 12.125±0.145 |




Fig. 1: Superimposed IR spectra of Stigmasterol (standard and isolated) in *C. setigerus*





Fig. 2: Mass Spectrum of Stigmasterol (RT: 34.342)



Graph 1: Comparative study of Antioxidants of *Cenchrus setigerus* (In-vitro and In-vivo)